



# Phytochemical analysis and hepatoprotective properties of *Tinospora cordifolia* against carbon tetrachloride-induced hepatic damage in rats

B. T. Kavitha<sup>1</sup>, S. D. Shruthi<sup>1, 3</sup>, S. Padmalatha Rai<sup>2</sup> and Y. L. Ramachandra<sup>1\*</sup>

<sup>1</sup>P.G. Department of Studies and Research in Biotechnology and Bioinformatics, Kuvempu University, Shankaraghatta – 577 451, Karnataka, India

<sup>2</sup>Department of Biotechnology, Manipal Life Science Center, Manipal University, Manipal – 576104, Karnataka, India

<sup>3</sup>P.G. Department of Biotechnology, The Oxford College of Science, Bangalore – 560 102, Karnataka, India

## ABSTRACT

The present study was conducted to evaluate the hepatoprotective activity of different extracts of *Tinospora cordifolia* against carbon tetrachloride (CCl<sub>4</sub>) induced liver damage in rats. The pet ether, ethanol and aqueous extracts of various parts of the plant such as leaf, stem and root were tested at the dose of 200mg/kg body weight orally using Wistar albino rats and Silymarin was given as reference standard. Ethanolic extract of all the parts showed significant hepatoprotective effect by reduction in serum enzymes alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase (ALP) and total bilirubin (TBL) in the selected model which is followed by aqueous and pet ether extracts. The chemical constituents reported from the plant belong to different classes such as alkaloids, flavonoids, glycosides, steroids, terpenoids, phenolics and saponins. The overall experimental results suggests that the biologically active phytoconstituents such as flavonoids, alkaloids present in the ethanolic extract of plant *Tinospora cordifolia*, may be responsible for the significant hepatoprotective activity. Therefore, results justify the use of *Tinospora cordifolia* as a hepatoprotective agent.

## KEYWORDS

Carbon tetrachloride, Extracts, Hepatoprotective activity, Phytochemical, *Tinospora cordifolia*.

received on 16-04-2011

accepted on 13-05-2011

available online 15-08-2011

www.jbclinpharm.com

## INTRODUCTION

*Tinospora cordifolia* (Guduchi) is a widely used plant in folk and ayurvedic systems of medicine. It is a climbing shrub belonging to the family Menispermaceae, which is widely distributed throughout Indian subcontinent and China [1]. The stem of *Tinospora cordifolia* is rather succulent with long filiform fleshy aerial roots from the branches. The bark is creamy white to grey, deeply left spirally, the space in between being spotted with large rosette like lenticels [2]. The stem is bitter, stomachic, diuretic [3], stimulates bile secretion, causes constipation, allays thirst, burning sensation, vomiting, enriches the blood and cures jaundice. The extract of its stem is useful in skin diseases [4] [5]. The root and stem of *T. cordifolia* are prescribed in combination with other drugs as an antidote to snake bite and scorpion sting [6] [7]. Oral administration of an aqueous *T. cordifolia* root extract to alloxan diabetic rats caused a significant reduction in blood glucose and brain lipids [8]. *T. cordifolia* is reported to benefit the immune system in a variety of ways [9]. The hepatoprotective action of *T. cordifolia* was reported in one of the experiment in which goats treated with *T. cordifolia* have shown significant clinical and hemato-biochemical improvement in CCl<sub>4</sub> induced hepatopathy. Extract of *T. cordifolia* has also exhibited *in vitro* inactivating property against Hepatitis B and E surface antigen [10]. The aqueous extract of *T. cordifolia* exerted a significant anti-inflammatory effect on cotton pellet granuloma and formalin induced arthritis models [11]. In a clinical evaluation, a compound preparation 'Rumalaya' containing *T. cordifolia* was reported to significantly reduce the pain in patients suffering from rheumatoid arthritis [12].

Liver diseases are a serious health problem. In the absence of reliable liver protective drugs in allopathic medical practices, herbs play a role in the management of various liver disorders. Numerous medicinal plants and their formulations are used

for liver disorders in ethnomedical practices and in traditional system of medicine in India [13]. In view of severe undesirable side effects of synthetic agents, there is growing focus to follow systematic research methodology and to evaluate scientific basis for the traditional herbal medicines that are claimed to possess hepatoprotective activity. The use of medicinal plants in modern medicine suffers from the fact that though hundreds of plants are used in the world to prevent or to cure diseases, scientific evidence in terms of modern medicine is lacking in most cases. However today it is necessary to provide scientific proof as whether to justify the use of plant or its active principles [14]. So, in the present study, we investigated the hepatoprotective effect of various parts of *T. cordifolia* extracted with pet ether, ethanol and aqueous against CCl<sub>4</sub> induced liver damage in rats.

## MATERIALS AND METHODS

### Plant materials and preparation of extracts

The different parts such as leaf, stem and root of the plant *Tinospora cordifolia* were freshly collected in winter season locally from Bhadra Wild Life Sanctuary, Karnataka (Southern India) in February 2010. The taxonomic identification of the plant was confirmed by Dr. Y. L. Ramachandra, Department of Biotechnology, Kuvempu University, Shankaraghatta (Voucher specimen number YLR530), has been preserved in Dept. of Biotechnology, Kuvempu University, Shankaraghatta for future references. The shade dried leaves were pulverised and subjected for extraction using pet ether, ethanol (LR grade, Merck, India) and aqueous separately using soxhlet apparatus. Then the extracts were concentrated using rotary flash evaporator (Buchi Flawil, Switzerland). The suspension of extracts were prepared using 5% gum acacia and used for the study.

### Preliminary phytochemical analysis

All the different extracts were then subjected to preliminary phytochemical analysis to assess the presence of various phytoconstituents [15].

\*Corresponding Author E-mail: ylpub@gmail.com

## Animal models

Studies were carried out using Wistar strain albino rats (150-200 g) of either sex, which were procured from Central Animal House, National College of Pharmacy, Shivamogga. They were housed under standard laboratory conditions (temp 23±2°C, relative humidity 55±10%) with dark light cycle (14/10hr). Animals were allowed free access to standard pellet diet (Sai Durga Feeds, Bangalore) and water ad libitum. All experimental protocols were prepared and performed based on ethical guidelines of Institutional Animal Ethics Committee (No. 144/1999/CPCSEA/SMG).

## Toxicity studies

Acute toxicity study was performed for all the extracts according to the staircase method of Ghosh [16]. Female albino rats were used for acute toxicity study. The animals were kept fasting for overnight providing only water, after which the extract was administered orally at doses of 100, 200 and 300mg/kg b.w. and observed for toxic symptoms up to 72 h. 200 mg/kg b.w was taken as the therapeutic oral dose for all the extracts.

## Carbon tetrachloride induced hepatotoxicity

Rats were divided into twelve groups of six animals each. The rats of control group (I) received three doses of 5% gum acacia mucilage (1ml/kg b.w.) at 12 h intervals (0 h, 12 h and 24 h). The rats of CCl<sub>4</sub> group (II) received three doses of vehicle at 12 h intervals and a single dose of CCl<sub>4</sub> (1.25ml/kg b.w.) diluted in liquid paraffin (1:1) 30 minutes after the administration of 1<sup>st</sup> dose of vehicle. Group – III served as reference control, received silymarin (25mg/kg b.w.) once daily for 3 days. Group –IV to XII received, *T. cordifolia* extract (200 mg/kg b.w.) once daily for 3 days. Group III to XII received CCl<sub>4</sub> (3gm/kg b.w.) as single dose on day 3, thirty minutes after the administration of extracts and silymarin respectively. All the test drugs and CCl<sub>4</sub> were administered orally by suspending in 5% gum acacia mucilage. After 36 hour of administration of CCl<sub>4</sub>, blood was collected by direct cardiac puncture under light ether anesthesia and serum was separated and used for determination of biochemical parameters such as alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase (ALP) and total bilirubin (TBL) [17].

## Histopathological studies

The animals were sacrificed and the abdomen was cut open to remove the liver. The liver was fixed in Bouin's solution (mixture of 75 ml of saturated picric acid, 25 ml of 40% formaldehyde and 5ml of glacial acetic acid) for 12h, and then embedded

in paraffin using conventional methods [18] and cut into 5µm thick sections and stained using haematoxylin-eosin dye and finally mounted in di-phenyl xylene. The sections were then observed under microscope for histopathological changes in liver architecture and their photomicrographs were taken.

## Statistical analysis

Statistical analysis of the data was done by one-way ANOVA, followed by student t-test using software ezANOVA ver. 0.98. The level of significance was fixed at P<0.05.

## RESULTS AND DISCUSSION

The preliminary phytochemical investigations show the presence of various secondary metabolites as tabulated in table 1. The same extracts have shown significant hepatoprotective activities at the concentration (200 mg/kg) tested, although its pure form remains to be studied. The results of biochemical parameters revealed the elevation of enzyme level in CCl<sub>4</sub> treated group, indicating that CCl<sub>4</sub> induces damage to the liver (Table 2). A significant reduction was observed in TBL, ALT, AST and ALP levels in the groups treated with silymarin and leaf extracts of *T. cordifolia*. But the groups which received other two extracts such as stem and root, as test drug at the dose of 200mg/kg showed a less significant decrease in the elevated levels of enzymes. The results obtained for biochemical parameters are comparable with silymarin, the standard hepatoprotective drug. Therefore, as like silymarin the leaf extracts have restored the altered level of enzymes significantly. The enzyme levels were almost restored to the normal. It was found that the extract decreased the CCl<sub>4</sub> induced elevated levels of the enzymes in group III to XII, indicating the production of structural integrity of hepatocytic cell membrane or regeneration of damaged liver cells by the extract. It was observed that the size of the liver was enlarged in CCl<sub>4</sub> intoxicated rats but it was normal in drug treated groups. A significant reduction in liver weight supports this finding.

Normally, AST and ALP are present in high concentration in liver. Due to hepatocyte necrosis or abnormal membrane permeability, these enzymes are released from the cells and their levels in the blood increases. ALT is a sensitive indicator of acute liver damage and elevation of this enzyme in non hepatic diseases is unusual. ALT is more selectively a paranchymal liver enzyme than AST [19]. Assessment of liver function can be made by estimating the activities of serum ALT, AST, ALP and TBL which are enzymes originally present higher concentration in cytoplasm. When there is hepatopathy, these enzymes leak into the blood stream in conformity with the extent of liver damage [20]. Bilirubin is one of the most useful clinical clues to the severity of necrosis and its accumulation is a measure of binding, conjugation and excretory capacity of hepatocyte. Decrease in serum

**Table 1: Qualitative analysis of *T. cordifolia* plant extracts**


Plant constituent	Leaf			Stem			Root		
	Pet. Ether ext.	Ethanol ext.	Aqueous ext.	Pet. Ether ext.	Ethanol ext.	Aqueous ext.	Pet. Ether ext.	Ethanol ext.	Aqueous ext.
Carbohydrates	–	+	+	–	+	+	–	+	+
Proteins	–	+	+	–	+	+	–	+	+
Tannins	–	+	+	–	+	+	–	+	+
Saponins	+	+	+	+	+	+	+	+	+
Triterpenoids	+	+	+	+	+	–	+	+	+
Flavonoids	+	+	+	+	+	+	+	+	+
Quinones	–	–	–	–	–	–	–	–	–
Sterols	+	–	–	+	–	–	+	–	–
Glycosides	+	+	+	+	+	+	+	+	+
Alkaloids	+	+	+	+	+	+	+	+	+

Phytochemical test: '+' - Present and '-' - Absent .

**Table 2:** Effect of leaf, stem and root extracts of *T. cordifolia* on CCl<sub>4</sub> induced hepatotoxicity in rats enzymes.

Group (n)	TBL (mg/dL) ± SE	ALT (IU/L) ± SE	AST (IU/L) ± SE	ALP (IU/L) ± SE
Control (I)	0.85 ± 0.02	202.50 ± 2.28	254.00 ± 2.50	300.83 ± 3.23
CCl <sub>4</sub> (II)	2.05 ± 0.03**	423.33 ± 3.17**	483.83 ± 2.39**	522.83 ± 2.23**
CCl <sub>4</sub> + S (III)	0.93 ± 0.03	194.83 ± 2.36	246.67 ± 2.16	291.83 ± 2.43
CCl <sub>4</sub> + PL (IV)	1.31 ± 0.03**	226.00 ± 1.79**	274.00 ± 2.28**	335.17 ± 2.26*
CCl <sub>4</sub> + EL (V)	0.95 ± 0.04**	204.33 ± 2.25**	253.83 ± 1.97**	303.67 ± 2.54**
CCl <sub>4</sub> + AL (VI)	1.11 ± 0.05**	213.67 ± 2.29**	265.33 ± 2.82**	329.67 ± 18.00**
CCl <sub>4</sub> + PS (VII)	1.46 ± 0.06	286.17 ± 2.10	334.67 ± 2.25	383.50 ± 2.47
CCl <sub>4</sub> + ES (VIII)	1.23 ± 0.05**	256.50 ± 2.22**	303.17 ± 2.48**	355.67 ± 2.06*
CCl <sub>4</sub> + AS (IX)	1.31 ± 0.02	275.17 ± 2.07	325.83 ± 2.75	381.33 ± 1.26
CCl <sub>4</sub> + PR (X)	1.47 ± 0.01	307.33 ± 1.91	355.17 ± 1.82	404.00 ± 2.38
CCl <sub>4</sub> + ER (XI)	1.28 ± 0.01**	285.83 ± 1.82**	334.50 ± 1.89*	384.00 ± 2.08**
CCl <sub>4</sub> + AR (XII)	1.32 ± 0.01	295.33 ± 1.93	344.83 ± 2.04	393.50 ± 2.19

PL-Pet ether ext. of leaf; EL-Ethanol ext. of leaf; AL-Aqueous ext. of leaf; PS- Pet ether ext. of stem; ES- Ethanol ext. of stem; AS- Aqueous ext. of stem; PR- Pet ether ext. of root; ER- Ethanol ext. of root; AR- Aqueous ext. of root. S- Silymarin, n = 6 animals in each group, The values are mean ± S.E. \*\* P<0.01 compared to standard.

**Figure 1:** Histopathological changes showing effect of Control (A); Positive control (B); Silymarin (C) and Drug treated (D).

bilirubin after treatment with the extract indicates the effectiveness of it in normal functional status of the liver.

Histopathological examination of the liver section of the rats treated with toxicant showed intense centrilobular necrosis and vacuolization. The rats treated with silymarin and extracts along with toxicant showed sign of protection against these toxicants to considerable extent as evident from formation of normal hepatic cords

and absence of necrosis and vacuoles (fig. 1). The preliminary phytochemical studies have revealed the presence of flavonoids in the extracts of *T. cordifolia*, which supports the fact that usually flavonoids show hepatoprotective activity [21]. So in the present study, the hepatoprotective effect of *T. cordifolia* may be due to its flavonoid content. CCl<sub>4</sub> induced animals significantly lost their body weight and showed reduced food consumption as compared to control group. Animals of drug

treated group and standard group showed a significant increase in body weight and food consumption comparably. These findings suggested that extract administered has significantly neutralized the toxic effects of  $\text{CCl}_4$  and helped in regeneration of hepatocytes [22]. In accordance with these results, it can be said that the leaf extracts of *T. cordifolia* exhibited a hepato protective effect against carbon tetrachloride induced hepatotoxicity.

## CONCLUSION

The plant is used in ayurvedic, "Rasayanas" to improve the immune system and the body resistance against infections. The Ayurveda literature reports that it can cause constipation, if taken regularly in high doses; it has no side effect and toxicity. Yet the safety and the potential indications in human beings have to be established using modern methods. These studies place this indigenous drug a novel candidate for bioprospection and drug development for the treatment of liver disorders where satisfactory cure managements are still not available. Efforts are in progress to isolate and characterize the active principle, which is responsible for the hepatoprotective efficacy of this valuable medicinal plant. Further studies are needed to explain the exact mechanism of action in neutralizing the toxic effects.

## REFERENCES

1. Singh SS, Pandey SC, Shrivastava S, et al. Chemistry and medicinal properties of *Tinospora cordifolia*. *Indian J Pharmacol.* 2003; 35: 83-9.
2. Kirtikar KR, Basu BD. *Indian Medicinal Plants*, Vol 1. 2nd ed. New Connaught Place, Dehra Dun; 1975.
3. Nayampalli SS, Ainapure SS, Samant BD, et al. A comparative study of diuretic effects of *Tinospora cordifolia* and hydrochloro-thiazide in rats and a preliminary phase I study in human volunteers. *J Postgrad Med.* 1988; 34: 233-6.
4. Aiyer KN, Kolamall M. *Pharmacognosy of Ayurvedic Drugs*, Series I. 1st ed. Trivendram: The Central Research Institute; 1963.
5. Raghunathan K, Mitra R. *Pharmacognosy of Indigenous Drugs*. New Delhi: Central Council for Research In Ayurveda & Siddh; 1982.
6. Nadkarni KM, Nadkarni AK. *Indian Materia Medica*, Vol 1. 3rd ed. Mumbai: M/S Popular Prakasan Pvt. Ltd; 1976.
7. Zhao TF, Wang X, Rimando AM, et al. Folkloric medicinal plants: *Tinospora sagittata* var. *cravaniana* and *Mahonia bealei*. *Planta Medica.* 1991; 57: 505.
8. Dhaliwal KS. Method and composition for treatment of diabetes. *US Patent* 5886029; 1999.
9. Kapil A, Sharma S. Immunopotentiating compounds from *Tinospora cordifolia*. *J Ethnopharmacol.* 1997; 58: 89-95.
10. Mehrotra R, Katiyar CK, Gupta AP. Hepatoprotective compositions and composition for treatment of conditions related to hepatitis B and E infection. *US Patent* 749296; 2000.
11. Jana U, Chattopadhyay RN, Shw BP. Preliminary studies on anti-inflammatory activity of *Zingiber officinale* Rosc., *Vitex negundo* Linn. and *Tinospora cordifolia* (Willid) Miers in albino rats. *Indian J Pharmacol.* 1999; 31: 232-3.
12. Anonymous. *Wealth of India: Raw materials*, Vol X. New Delhi: CSIR; 1976.
13. Sethuraman MG, Lalitha KG, Raj Kapoor B. Hepatoprotective activity of *Sarcostemma brevistigma* against carbon tetrachloride-induced hepatic damage in rats. *Current Science.* 2003; 84: 1186-87.
14. Singh RP, Padmavathi B, Rao AR. Modulatory influence of *Adhatoda vesica* (*Justica adhatoda*) leaf extract on the enzyme of xenobiotic metabolism, antioxidant status and lipid peroxidation in mice. *Mol Cell Biochem.* 2000; 213: 99-109.
15. Harbone JB. *Phytochemical methods - A guide to modern technique of plant analysis*, 2nd edn, Chapman and Hall, New York; 1984. 85pp.
16. Ghosh MN. *Fundamentals of experimental pharmacology*. Scientific book agency, Calcutta; 1984. 156-157pp.
17. Rekha Rajendran, Hemalatha S, Akasakalai K, et al. Hepatoprotective activity of *Mimosa pudica* leaves against Carbon tetrachloride induced toxicity. *J Nat Prod.* 2009; 2: 116-122.
18. Galighor AE, Kozloff EN. *Essentials of practical microtechnique* 2nd edn, Lea and Febiger, New York; 1976.
19. Shah M, Patel P, Phadke M, et al. Evaluation of the effect of aqueous extract from powders of root, stem, leaves and whole plant of *Phyllanthus debilis* against  $\text{CCl}_4$  induced rat liver dysfunction. *Indian Drugs.* 2002; 39: 333-337.
20. Nkosi CZ, Opoku AR, Terblanche SE. Effect of pumpkin seed (*Cucurbita pepo*) protein isolate on the activity levels of certain plasma enzymes in  $\text{CCl}_4$ -induced liver injury in low protein fed rats. *Phytother Res.* 2005; 19: 341-345.
21. Scevola D, Baebacini GM, Grosso A, et al. Flavonoids and hepatic cyclic monophosphates in liver injury. *Bollettino Dell Istituto Sieroterapico Milanese.* 1984; 63: 77-82.
22. Farooq S, Ahmed I, Pathak GK. In protective role of koflet (an Ayurvedic preparation) against cellular toxicity caused by Carbon tetrachloride and fly-ash. *J Ethnopharmacol.* 1997; 53: 109-116.